

DAVID GRAYS TRICHLORPHON 60 Caterpillar and White Butterfly Spray.
SAFETY DIRECTIONS: The concentrate is poisonous. Avoid contact with skin and eyes. Do not inhale spray mist. When using the product, wear rubber gloves. Wash hands after use.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. If swallowed induce vomiting preferably using Ipecac Syrup. APF and give one atropine tablet every 15 minutes until dryness of mouth occurs. If poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly.
FLAMMABLE: KEEP AWAY FROM NAKED FLAME

POISON

**NOT TO BE TAKEN
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING**


**DAVID GRAYS
TRICHLORPHON 60
CATERPILLAR AND
WHITE BUTTERFLY**

**ACTIVE CONSTITUENT
600g/l TRICHLORPHON
(an anticholinesterase
compound)
200ml**

Packed in W.A. by
DAVID GRAY & CO., PTY. LIMITED
Rawlinson Street, O'Connor, W.A.

DAVID GRAYS TRICHLORPHON 60 Caterpillar and White Butterfly Spray
DIRECTIONS FOR USE: To control White Butterfly, Cabbage Moth, Green Vegetable Bug on vegetable crops or ornamentals, mix 12ml with 10 litres of water. Spray immediately on infested plants when pests first appear. Repeat weekly. To control Fruit Fly on Stone Fruit mix 20ml with 10 litres of water. Spray thoroughly 6 weeks before fruit ripens. Repeat weekly at half strength - last spray ONE WEEK before fruit ripens. To control Fruit Fly on Stone Fruit mix 40ml with 10 litres for first spray, then apply as for Stone Fruit.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER
CONTRARY TO THIS LABEL UNLESS AUTHORISED.
DO NOT APPLY LATER THAN 2 DAYS BEFORE HARVESTING OR GRAZING.



POISON

NOT TO BE TAKEN

KEEP OUT OF REACH OF CHILDREN


READ **SAFETY** DIRECTIONS BEFORE OPENING

DAVID GRAYS

TRICHLORPHON 60

**CATERPILLAR
& WHITE
BUTTERFLY
SPRAY**

ACTIVE CONSTITUENT
600g/l TRICHLORPHON



200 ml

**DAVID GRAYS CATERPILLAR
& WHITE BUTTERFLY SPRAY**

SAFETY DIRECTIONS:

The concentrate is poisonous. Avoid contact with skin and eyes and avoid breathing vapour. When handling the concentrate and preparing spray use rubber gloves and face shield. If spilled on skin and on completion of spraying wash thoroughly with soap and water. Wash contaminated clothing before reuse. Do not eat or smoke while spraying. Store in original container, tightly closed in a safe place. Wash out container thoroughly and dispose of safely.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. If swallowed induce vomiting, Use Ipecac Syrup (APF) if available. Give one Atropine tablet every quarter hour until dryness of mouth occurs. If poisoned by skin absorption remove contaminated clothing and wash skin thoroughly. Give Atropine tablets as above.



9 310299 078453

**SPRAY
BUTTERFLY
& WHITE
CATERPILLAR**

TRICHLORPHON 60

DAVID GRAYS



POISON

NOT TO BE TAKEN

KEEP OUT OF REACH OF CHILDREN


READ SAFETY DIRECTIONS BEFORE OPENING

DAVID GRAYS

TRICHLORPHON 60

**CATERPILLAR
& WHITE
BUTTERFLY
SPRAY**

ACTIVE CONSTITUENT
600g/l TRICHLORPHON



**DAVID GRAYS CATERPILLAR
& WHITE BUTTERFLY SPRAY**

DIRECTIONS FOR USE:

To control White Butterfly, Cabbage Moth, Green Vegetable Bug on vegetable crops or ornamentals, mix 12ml with 10 litres of water. Spray immediately on infested plants when pests first appear.

Repeat weekly.

To control Fruit Fly on STONE FRUIT mix 20ml with 10 litres of water. Spray thoroughly 6 weeks before fruit ripens. Repeat every week at half strength, last spray ONE WEEK before fruit ripens. On POME FRUIT mix 40ml in 10 litres of water for first spray, then apply as for stone fruit.

NOT TO BE USED FOR ANY PURPOSE OR ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.

Do NOT apply later than 2 days before harvesting.



Formulated and Packed by
DAVID GRAY & CO. PTY. LIMITED
RAWLINSON ST., O'CONNOR W.A.